



Dental Clinical Policy

Subject: Anesthesia

Guideline #: 09-201

Status: Revised

Publish Date:

01/01/2026

Last Review Date:

10/27/2025

Description

The services identified in this policy are a part of a continuum of depth of sedation that is widely accepted and sets the standards for defining levels of sedation. The level of anesthesia is dependent upon the effects of the anesthetic agent on the central nervous system.

Deep sedation/general anesthesia is a drug-induced loss of consciousness where patients are unarousable, even by painful stimulation. In this state, cardiovascular function might also be impaired. No interventions are necessary to maintain an airway, spontaneous breathing is adequate, and cardiovascular function is usually maintained.

Intravenous moderate sedation is a level of drug induced depression of consciousness used to help a person feel relaxed and calm. No interventions are necessary to maintain an airway, spontaneous breathing is adequate, and cardiovascular function is usually maintained. The person remains awake, able to follow directions and typically remembers little to nothing of the procedure completed.

Analgesia refers to a drug-induced depression of consciousness in which a person is not easily aroused but responds purposefully after repeated or painful stimulation. Assistance may be required to maintain an airway and spontaneous breathing may be inadequate. Cardiovascular function is generally preserved.

Anxiolysis is a drug induced state of consciousness which allows a person to respond normally to verbal commands. While cognitive function and coordination may be impaired, a person maintains their airway reflexes, breathing and cardiovascular function.

Clinical Indications for Anesthesia Services

The administration of local anesthesia, nitrous oxide, intravenous moderate sedation and deep sedation/general anesthesia is an integral part of dental practice. Dental professionals administering anesthesia must be appropriately educated, trained, and licensed.

The American Dental Association recommends avoidance of the use of preoperative oral sedatives for children (ages 12 and under) prior to arrival in the dental office, except in extraordinary situations due to the risk of unobserved respiratory obstruction during transport to the dental office by untrained individuals. For children 12 years of age and under, the American Dental Association supports the use of the American Academy of Pediatrics/American Academy of Pediatric Dentistry Guidelines for Monitoring and Management of Pediatric Patients During and After Sedation for Diagnostic and Therapeutic Procedures.

According to The American Academy of Pediatrics (AAP) and the American Academy of Pediatric

Dentistry (AAPD), the definition of a child is between the ages of 0 to 21. While this definition may be appropriate to define a child by these recognized academies, for benefit purposes, Anthem follows criteria and state mandated ages for Intravenous (IV) and inhalation anesthesia for children.

When more advanced anesthetics, such as intravenous (IV) or inhalation are used to induce more profound levels of anesthesia, additional individuals trained in either Advanced or Basic Life Support (state specific) for Healthcare Providers must be present in addition to the dentist/oral surgeon. If required by the State, a current equipment inspection certificate, sedation license and CPR certification are mandatory.

Deep sedation/general anesthesia or intravenous moderate sedation may be considered when performed in conjunction with dental procedures other than oral surgery when deemed necessary due to concurrent individual medical conditions/diagnoses. These conditions may be inclusive of, but not limited to, physical, intellectual or medically compromised conditions for which dental treatment using a local anesthetic cannot be expected to provide a successful result. Medical conditions, which may necessitate general anesthesia or moderate sedation services, must be supported by medical documentation from a physician.

Medical Necessity and Generally Accepted Standards of Care

Medically/Dentally Necessary or Medical/Dental Necessity means Medical/Dental Services that are:

(1) Consistent with the Member's diagnosis or condition.

(2) Rendered:

- In response to a life-threatening condition or pain; or
- To treat an injury, illness or infection related to the dentition; or
- To achieve a level of function to the dentition consistent with prevailing community standards for the diagnosis or condition.

Monitoring and Documentation:

1. Monitoring: A dentist, or at the dentist's direction, an appropriately trained individual, must remain in the operatory during active dental treatment to monitor the patient's vital signs continuously until the patient meets the criteria for discharge to the recovery area. The appropriately trained individual must be familiar with monitoring techniques and equipment. Monitoring must include observation of proper oxygenation (includes observation of the color of mucosa – pink hue); skin or blood must be continually evaluated. Oxygen saturation by pulse oximetry is appropriate and necessary.
2. Ventilation: chest movement must be continually observed. Respirations must be verified and documented by appropriately trained personnel.
3. Circulation: Blood pressure and heart rate must be evaluated pre-operatively, postoperatively and intra-operatively and properly documented.
4. Documentation: An appropriate, complete anesthesia record must be maintained that includes the names of all drugs administered, time of administration including dosages of local anesthetics or other anesthetic drugs (IV or inhalation). All physiological parameters must be monitored and properly documented.

Emergency Management:

The dentist is responsible for drug management, adequacy of the facility and staff, diagnosis and treatment of emergencies related to the administration of intravenous or inhalation drugs as well as providing the appropriate equipment, drugs and office protocol for emergency management.

Criteria

Local Anesthesia

- The administration of local anesthetic; be it traditional, electronic or buffered etc. - is

considered inclusive of (part of) all dental procedures [unless a specific plan allows coverage] and is not eligible for a separate benefit.

Regional and Trigeminal Division Block Anesthesia

- Regional and trigeminal block anesthesia may not be a covered service.

Deep Sedation/General Anesthesia, Intravenous Moderate Sedation/Analgesia, Non- Intravenous Conscious Sedation

- These procedures may be benefitted when appropriate in conjunction with covered dental services.
- Medical conditions, which require the use of general anesthesia, must be supported by documentation submitted by the oral surgeon, dentist and/or physician administering the anesthesia.
- To qualify for general anesthesia or moderate sedation benefit, the member must satisfy one of the criteria noted below.
 - a The member is a child, up to the age defined by contract, with a dental condition (such as Early Childhood Caries) that requires repairs of significant complexity (for example, multiple amalgam and/or resin based composite restorations, pulpal therapy, extractions or any combinations of procedures as noted or other dental procedures)
 - b The member has physical, intellectual, or medically compromising conditions for which dental treatment under local anesthesia, with or without additional adjunctive techniques and modalities, cannot be expected to provide a successful result and which, under anesthesia, can be expected to produce a superior result. Conditions include but are not limited to intellectual disabilities, cerebral palsy, epilepsy, severe cardiac problems and hyperactivity (verified by appropriate medical documentation)
 - c The member is extremely uncooperative, fearful, unmanageable, anxious, or uncommunicative with complex dental needs which should not be postponed or deferred and for whom lack of treatment can be expected to result in oral pain, infection, loss of teeth, or other increased oral morbidity.
 - d The member for whom local anesthesia is ineffective (for reasons such as acute infection, anatomic variations or allergy). Failed attempts of local anesthesia administration must be documented and submitted for review
 - e The member has sustained extensive oral-facial and/or dental trauma, for which treatment under local anesthesia would be ineffective or compromised
- The level of anesthesia is not determined by the route of administration, but by the dentist's documentation of the anesthetic agent's effect on the patient's central nervous system.
- The time of general anesthesia begins when the dentist starts administering the sedative agent and non-invasive monitoring protocol.
- The dentist administering the anesthesia must remain in constant attendance with the patient until completion of the anesthesia procedure.
- Services are considered complete when the patient may be safely left under the observation of trained personnel, and the dentist may safely leave the operatory to attend to other patients.
- Office anesthesia in excess of 60 minutes for any dental or surgical procedure requires

written rationale/documentation explaining the necessity. Necessary documentation includes all associated radiographic images, progress notes, operative report and a complete anesthesia record indicating start and stop times of incremental drug administration.

Inhalation of Nitrous Oxide/Analgesia

- These procedures may be benefitted when appropriate in conjunction with covered dental services.
- Nitrous Oxide may be indicated for:
 - a. ineffective local anesthesia administration
 - b. treatment of dental anxiety
 - c. special needs patients
 - d. individuals who are uncooperative or behaviorally challenged
- Nitrous Oxide is contraindicated for, but not limited to:
 - a. Patients with severe respiratory compromise
 - b. First trimester pregnancy because of its potential teratogenic effects
 - c. Patients with a history of stroke
 - d. Hypotension and known cardiac conditions

Coding

The following codes for treatments and procedures applicable to this document are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

CDT including but not limited to:

- D9210 local anesthesia not in conjunction with operative or surgical procedures
- D9211 regional block anesthesia
- D9212 trigeminal division block anesthesia
- D9215 local anesthesia in conjunction with operative or surgical procedures
- D9219 evaluation for deep sedation or general anesthesia
- D9222 administration of deep sedation/general anesthesia – first 15-minute increment, or any portion thereof
- D9223 administration of deep sedation/general anesthesia – each subsequent 15-minute increment, or any portion thereof
- D9224 administration of general anesthesia with advanced airway – first 15-minute increment, or any portion thereof
- D9225 administration of general anesthesia with advanced airway – each subsequent 15-minute increment, or any portion thereof
- D9230 administration of nitrous oxide administration of moderate sedation – intravenous
- D9239 – first 15-minute increment, or any portion thereof
- D9243 administration of moderate sedation – intravenous – each subsequent 15-minute increment, or any portion thereof
- in-office administration of minimal sedation – single drug – enteral
- D9244 administration of moderate sedation – enteral
- D9245 administration of moderate sedation – non-intravenous parenteral – first 15-minute increment, or any portion thereof
- administration of moderate sedation – non-intravenous parenteral – each

D9247 subsequent 15-minute increment, or any portion thereof

D9613 infiltration of sustained release therapeutic drug, per quadrant

ICD-10 CM Diagnoses for Dental Diseases and Conditions: See the current CDT code book for details

References

1. American Society of Anesthesiologists. Statement on Continuum of Depth of Sedation: Definition of General Anesthesia and Levels of Sedation/Analgesia. asahq.org. Last Amended: October 23, 2019.
2. American Academy of Pediatric Dentistry. Behavior Guidance for the Pediatric Dental Patient. aapd.org. Updated 2020.
3. American Academy of Pediatric Dentistry. Policy on Patient Safety. aapd.org. Updated 2021.
4. American Academy of Pediatric Dentistry. Use of Nitrous Oxide for Pediatric Dental Patients. aapd.org. Updated 2018.
5. American Academy of Pediatric Dentistry. Guidelines for Monitoring and Management of Pediatric Patients Before, During, and After Sedation for Diagnostic and Therapeutic Procedures. aapd.org. Published August 19, 2019.
6. American Society of Anesthesiologists. Statement on Sedation & Anesthesia Administration in Dental Office-Based Settings*. asahq.org. Updated October 26, 2022.
7. American Dental Association. Guidelines for the Use of Sedation and General Anesthesia by Dentists.ada.org. Adopted October 2016.
8. CDT 2026 Current Dental Terminology, American Dental Association.

History

Revision History	Version	Date	Nature of Change	SME
	Initial	01/01/2014		Dr. Koumaras
	Revision	02/08/2017		
	Revision	02/06/2018	Appropriateness and medical necessity	Dr. Kahn
	Revised	11/12/2020	Annual Review	Committee
	Revised	12/06/2020	Annual Review	Committee
	Revised	10/20/2021	Annual Review	Committee
	Revised	11/11/2022	Annual Review	Committee
	Revised	11/01/2023	Annual Review	Committee
	Revised	10/31/2023	Minor editorial refinements to description, clinical indications, criteria, and reference; intent unchanged.	Committee
	Reviewed	10/27/2025	Minor editorial refinements to description, clinical	Dr. Balikov

		indications, criteria, and reference; intent unchanged.	
--	--	---	--

Federal and State law, as well as contract language, takes precedence over Dental Clinical Policy. Dental Clinical Policy provides guidance in interpreting dental benefit applications. The Plan reserves the right to modify its Dental Clinical Policies and guidelines periodically and as necessary. Dental Clinical Policy is provided for informational purposes and does not constitute medical advice. These Policies are available for general adoption by any lines of business for consistent review of the medical or dental necessity and/or appropriateness of care of dental services. To determine if a review is required, please contact the customer service number on the member's card.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the Plan.

Current Dental Terminology - CDT © 2026 American Dental Association. All rights reserved.